arXiv:math/0603467v2 [math.GT] 19 Sep 2008

QUANTUM HYPERBOLIC INVARIANTS FOR DIFFEOMORPHISMS OF SMALL SURFACES

XIAOBO LIU

Abstract. An earlier article [1] introduced new invariants for pseudo-Anosov diffeomorphisms of surface, based on the representation theory of the quantum Teichmüller space. We explicity compute these quantum hyperbolic invariants in the case of the 1–puncture torus and the 4–puncture sphere.

1. Introduction

In [1], Francis Bonahon and the author constructed quantum hyperbolic invariants for pseudo-Anosov diffeomorphisms of a punctured surface S. For every odd integer N, these invariants associate to the pseudo-Anosov diffeomorphism $\varphi : S \to S$ a square matrix C_φ of dimension N^{3g+p-3} (where $g \geq 0$ is the genus of S and $p > 0$ is its number of punctures), defined up to conjugation and scalar multiplication. In particular, C_φ is an $N \times N$ matrix for the 1–puncture torus and the 4–puncture sphere. The current paper provides an explicit computation for these two surfaces. It is based on the fact that the mapping class group of these surfaces is particularly simple.

For the 1-puncture torus, the isotopy class of a diffeomorphism $\varphi : S \to S$ is completely determined by the matrix $A_\varphi \in \mathrm{SL}_2(\mathbb{Z})$ defined by considering the action of φ on $H_1(S) = \mathbb{Z}^2$. A diffeomorphism φ is pseudo-Anosov if and only if $|\mathrm{Tr}(A_\varphi)| > 2$. Such a matrix A_φ is conjugate to a product

$$A_\varphi = A_1 A_2 \cdots A_n, \quad \text{where} \quad A_i = R = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \text{ or } A_i = L = \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}.$$

Our computation makes use of the matrices $C_R(u, v, u', v', h)$ and $C_L(u, v, u', v', h)$ defined by

$$\begin{aligned} &C_R(u, v, u', v', h)_{ij} \\ &= q^{2(j-i)^2} \left(\frac{u'v'}{uh}\right)^{j-i} \left(\frac{v'}{v}\right)^{i} \prod_{\alpha=1}^{i} \frac{1}{\left(1 + q^{4\alpha-3}u\right)\left(1 + q^{4\alpha-1}u\right)}, \end{aligned}$$

and

$$\begin{aligned} &C_L(u, v, u'', v'', h)_{ij} \\ &= \sum_{k=1}^{N} q^{4ik}\, q^{2(j-k)^2+2k^2} \left(\frac{u''v''}{vh}\right)^{j-k} \left(\frac{uvv''}{h}\right)^{k} \prod_{\alpha=1}^{k} \frac{1}{\left(1 + q^{4\alpha-3}v\right)\left(1 + q^{4\alpha-1}v\right)}. \end{aligned}$$

Theorem 1. *Let $\varphi : S \to S$ be a pseudo-Anosov diffeomorphism of the 1–puncture torus S, and consider a matrix $A_\varphi = A_1 A_2 \cdots A_n$, with $A_i = R$ or L, associated to φ as above. Then*

$$C_\varphi = C_1 C_2 \cdots C_n$$

where $C_i = C_R(u_{i-1}, v_{i-1}, u_i, v_i, 1)$ if $A_i = R$, $C_i = C_L(u_{i-1}, v_{i-1}, u_i, v_i, 1)$ if $A_i = L$, and where the complex numbers u_i, v_i are explicitly defined in §4 and §5 and are determined by the complete hyperbolic metric of the mapping torus M_φ of φ.

There is a similar theorem in the case of the 4-puncture sphere, using different matrix functions C_R^* and C_L^* defined in §6. In this situation, the action on slopes defines a map from the mapping class group of

---

Date: September 19, 2008.

1991 *Mathematics Subject Classification.* Primary 57R56; Secondary 57M50, 20G42.

Key words and phrases. mapping torus, representations.

This work was partially supported by NSF grant DMS-0103511 at the University of Southern California.

the 4-puncture sphere to $\mathrm{PSL}_2(\mathbb{Z})$ with kernel $\mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}/2\mathbb{Z}$. The image of a diffeomorphism φ can be represented by a matrix $A_\varphi \in \mathrm{SL}_2(\mathbb{Z})$. Again, φ is pseudo-Anosov if and only if $|\mathrm{Tr}(A_\varphi)| > 2$.

Theorem 2. *Let $\varphi : S \to S$ be a pseudo-Anosov diffeomorphism of the 4–puncture sphere S, and consider a matrix $A_\varphi = A_1 A_2 \cdots A_n$, with $A_i = R$ or L, associated to φ as above. Then*

$$C_\varphi = C_1^* C_2^* \cdots C_n^*$$

where $C_i^* = C_R^*(u_{i-1}, v_{i-1}, u_i, v_i, 1)$ if $A_i = R$, $C_i^* = C_L^*(u_{i-1}, v_{i-1}, u_i, v_i, 1)$ if $A_i = L$, and where the complex numbers u_i, v_i are explicitly defined in §6 and are determined by the complete hyperbolic metric of the mapping torus M_φ of φ.

Acknowledgements: We would like to thank François Guéritaud for crucial help with the combinatorics of the 1–puncture torus, and Francis Bonahon for general advice and encouragement.

2. Quantum hyperbolic invariants of surface diffeomorphisms

We briefly sketch the construction in [1] of quantum hyperbolic invariants for pseudo-Anosov diffeomorphisms of a punctured surface. Details will be provided in later sections in the specific case of the 1–puncture torus and the 4–puncture sphere.

Let S be a punctured surface, that is, a closed surface with finitely many points removed. The Teichmüller space of S, denoted by $\mathcal{T}_S$, is the space of isotopy classes of complete hyperbolic metrics on S. An ideal triangulation λ provides a certain set of global coordinates for $\mathcal{T}_S$, called the exponential shear coordinates (these essentially are cross-ratios), which parameterize this space by a convex cell $\mathbb{R}_+^n$ (more accurately they parametrize a finite-sheeted branched covering of $\mathcal{T}_S$ called the enhanced Teichmüller space).

The *quantum Teichmüller space* of S is a noncommutative deformation of the algebra of rational functions on $\mathcal{T}_S$, with a deformation parameter $q \in \mathbb{C}$; see [2, 4, 1, 5]. The construction is based on a non-commutative analog of the shear coordinate parametrization for the quantum Teichmüller space, the *Chekhov-Fock algebra* $\mathcal{T}_\lambda^q$ associated to an ideal triangulation λ of S.

When we change the ideal triangulation λ to another one λ', the exponential shear coordinates transform rationally. There is a non-commutative procedure that mimics this transformation rule of cross-ratios: it consists of an algebraic isomorphism $\Phi_{\lambda\lambda'}^q : \widehat{\mathcal{T}}_{\lambda'}^q \to \widehat{\mathcal{T}}_\lambda^q$, where $\widehat{\mathcal{T}}_\lambda^q$ is the division algebra of formal fractions of $\mathcal{T}_\lambda^q$. This *quantum coordinate change isomorphism* was originally proposed by Chekhov and Fock, and expressed in terms of a holomorphic function called the quantum dilogarithm (see [2]).

The Chekhov-Fock algebra $\mathcal{T}_\lambda^q$ admits finite dimensional representations only when q is a root of unity. Assume that q is a primitive N–root of unity with N odd. Any finite dimensional irreducible representation of this algebra has dimension N^{3g+p-3} (where $g \geq 0$ is the genus of S and $p > 0$ is its number of punctures) and, up to finitely many choices, is determined by complex weights $x_e \in \mathbb{C} - \{0\}$ associated to the edges e of the ideal triangulation λ.

It turns out that this representation theory of the Chekhov-Fock algebras $\widehat{\mathcal{T}}_\lambda^q$ is well-behaved with respect to the coordinate change isomorphisms $\Phi_{\lambda\lambda'}^q$. Namely, suppose that the irreducible representation $\rho : \mathcal{T}_\lambda^q \to \mathrm{End}(V)$ is associated to weights x_e, as e ranges over all edges of λ, and that these weights are in the domain of the (complexification of the) crossratio transformation rule between the exponential shear coordinates associated to λ and those associated to λ'. Then it is possible to make sense of the irreducible representation $\rho' = \rho \circ \Phi_{\lambda\lambda'}^q : \mathcal{T}_{\lambda'}^q \to \mathrm{End}(V)$, and ρ' is classified by the edge weights on λ' that are the image of the weights x_e under the cross-ratio transformation rule.

Now, consider a pseudo-Anosov diffeomorphism $\varphi : S \to S$. Thurston's Hyperbolization Theorem and Mostow's Rigidity Theorem provide a unique finite-volume complete hyperbolic metric on the mapping torus $M_\varphi = S \times [0,1]/\sim$, where $\sim$ identifies each $(x, 0)$ with $(\varphi(x), 1)$. An arbitrary ideal triangulation λ determines a unique pleated surface $f_\lambda : S \to M_\varphi$ with pleating locus λ, namely such that f sends each face of λ to a totally geodesic triangle in M_φ. This pleated surface defines a complex weight $x_e \in \mathbb{C} - \{0\}$ for each edge e of λ, called its exponential shear-bend parameter. If we identify λ and $\lambda' = \varphi(\lambda)$, these shear-bend coordinates have the property that they are invariant under the cross-ratio transformation rule from λ to λ'. Using the correspondence between edge weights and representations of the Chekhov-Fock algebra, this gives a representation $\rho : \mathcal{T}_\lambda^q \to \mathrm{End}(V)$ such that $\rho' = \rho \circ \Phi_{\lambda\lambda'}^q$ is isomorphic to $\rho \circ \Phi$, using the identification

$\Phi : \mathcal{T}^q_{\lambda'} \cong \mathcal{T}^q_{\lambda}$ provided by φ. In other words, there exists an isomorphism $C_\varphi \in \mathrm{GL}(V)$ such that

$$\rho \circ \Phi^q_{\lambda\varphi(\lambda)}(X) = C_\varphi \cdot \Big(\rho \circ \Phi(X)\Big) \cdot C_\varphi^{-1} \qquad \text{for any} \quad X \in \mathcal{T}^q_{\varphi(\lambda)}. \tag{2.1}$$

Theorem 3 (Theorem 40 in [1])**.** *The isomorphism $C_\varphi \in \mathrm{GL}(V)$ defined above depends only on q and φ, up to conjugation and scalar multiplication.* □

This article is devoted to computing the invariant C_φ when the surface S is the 1–puncture torus or the 4-puncture sphere. In these cases, the irreducible representations of the Chekhov-Fock algebra have dimension N, so that C_φ can be considered as an $N \times N$ matrix defined up to conjugation and scalar multiplication.

3. Quantum Teichmüller space of the 1–puncture torus

Following the construction of the quantum enhanced Teichmüller space [5], fix an *ideal triangulation* λ of the 1-puncture torus T, namely a triangulation of the unpunctured torus $\bar{T}$ with a single vertex, located at the puncture. The *Chekhov-Fock algebra* $\mathcal{T}^q_\lambda$ is defined by generators $X_1^{\pm 1}$, $X_2^{\pm 1}$, and $X_3^{\pm 1}$, respectively associated to the edges λ_1, λ_2, λ_3 of λ, and by relations determined by the topology of λ as follows:

$$X_i X_j = q^{2\sigma_{ij}} X_j X_i$$

where $\sigma_{ij} \in \{-2,-1,0,1,2\}$ is the number of times one sees $\lambda_i\bigwedge\lambda_j$ minus the number of times one sees $\lambda_j\bigwedge\lambda_i$. In particular, the σ_{ij} are antisymmetric in the subscripts. When the λ_i occur counterclockwise around the faces of λ as in Figure 1, then $\sigma_{ij} = -2$ whenever $j = i+1 \mod 3$.



Figure 1. An ideal triangulation λ. In this ideal triangulation $\sigma_{12} = \sigma_{23} = \sigma_{31} = -2$.

The *diagonal exchange* Δ_3 takes the ideal triangulation λ to a new ideal triangulation λ' obtained by replacing λ_3 by the other diagonal λ_3' of the square formed by λ_1 and λ_2, as Figure 2. If $\widehat{\mathcal{T}}^q_\lambda$ denotes the fraction division algebra of $\mathcal{T}^q_\lambda$, consisting of rational functions in the skew-commuting variable X_1, X_2, X_3, the coordinate change isomorphism $\Phi^q_{\lambda\lambda'} : \widehat{\mathcal{T}}^q_{\lambda'} \to \widehat{\mathcal{T}}^q_{\lambda}$ introduced in [5] is such that

$$\begin{aligned} \Phi^q_{\lambda\lambda'}(X_1') &= (1+qX_3)(1+q^3X_3)X_1, \\ \Phi^q_{\lambda\lambda'}(X_2') &= \big(1+qX_3^{-1}\big)^{-1}\big(1+q^3X_3^{-1}\big)^{-1}X_2, \\ \Phi^q_{\lambda\lambda'}(X_3') &= X_3^{-1}. \end{aligned} \tag{3.1}$$

in the case of Figure 1, namely if $\sigma_{ij} = -2$ when $j = i+1 \mod 3$.



Figure 2. The diagonal exchange Δ_3 on edge λ_3.

Now suppose that q is a primitive N-th root of unity with N odd. The center of $\mathcal{T}^q_\lambda$ is generated by X_1^N, X_2^N, and $X_1X_2X_3$.

Theorem 4 (A special case of Theorem 21 in [1])**.** *Let q be a primitive N-th root of unity with N odd. Any irreducible representation ρ of the Chekhov-Fock algebra is of dimension N, and the conjugacy class of ρ is determined by its restriction to the central elements X_1^N, X_2^N and $H = q^{-(\sigma_{12}+\sigma_{13}+\sigma_{23})}X_1X_2X_3$.* □

The coefficient introduced in the definition $H = q^{-(\sigma_{12}+\sigma_{13}+\sigma_{23})}X_1X_2X_3$ is designed to make H independent of the order of the λ_i. This will also guarantee that H is invariant under coordinate change isomorphisms in the sense that $\Phi^q_{\lambda\lambda'}(H') = H$ as proved in [5]. Also, note that $\sigma_{12}+\sigma_{13}+\sigma_{23} = \pm 2$, which may make this exponent less intimidating.

If $\rho : \mathcal{T}^q_\lambda \to \mathrm{End}(V)$ is an irreducible representation, then $\rho(X_1^N) = x_1 \,\mathrm{id}_V$, $\rho(X_2^N) = x_2 \,\mathrm{id}_V$, and $\rho(H) = h \,\mathrm{id}_V$ for complex numbers $x_1, x_2, h \in \mathbb{C} - \{0\}$. It follows that $\rho(X_3^N) = x_3 \,\mathrm{id}_V$ where $x_3 = h^N/(x_1x_2)$. In the next section we shall interpret (x_1, x_2, x_3) as the shear-bend coordinates of a pleated 1-puncture torus. The following lemma provides a base for this interpretation. Letting $q = 1$ in Formula (3.1) so that the variables commute, the isomorphism $\Phi^1_{\lambda\lambda'}$ defined there can be seen as a rational function from $\big(\mathbb{C} - \{0, -1\}\big)^3$ to itself. This $\Phi^1_{\lambda\lambda'}$ is exactly the transformation rule of the shear-bend coordinates of a pleated torus under diagonal exchanges.

Lemma 5 (Lemma 27 in [1])**.** *Let q be a primitive N-th root of unity with N odd, and let $\rho : \mathcal{T}^q_\lambda \to \mathrm{End}(V)$ be an irreducible representation with*

$$\rho(X_1^N) = x_1 \,\mathrm{id}_V, \quad \rho(X_2^N) = x_2 \,\mathrm{id}_V, \quad \rho(X_3^N) = x_3 \,\mathrm{id}_V, \quad \rho(H) = h \,\mathrm{id}_V$$

then $\rho' := \rho \circ \Phi^q_{\lambda\lambda'} : \mathcal{T}^q_{\lambda'} \to \mathrm{End}(V)$ is an irreducible representation with

$$\rho'\big({X'_1}^N\big) = x'_1 \,\mathrm{id}_V, \quad \rho'\big({X'_2}^N\big) = x'_2 \,\mathrm{id}_V, \quad \rho'\big({X'_3}^N\big) = x'_3 \,\mathrm{id}_V, \quad \rho(H') = h' \,\mathrm{id}_V$$

where $(x'_1, x'_2, x'_3) = \Phi^1_{\lambda\lambda'}(x_1, x_2, x_3)$ and $h' = h$. □

Note: Since $\Phi^q_{\lambda\lambda'}(\mathcal{T}^q_{\lambda'}) \nsubseteq \mathcal{T}^q_\lambda$, one needs Lemma 25 and Lemma 26 in [1] to rigorously make sense of $\rho \circ \Phi^q_{\lambda\lambda'}$ as a representation of $\mathcal{T}^q_{\lambda'}$. However, this turns out to coincide with our intuitive understanding.

4. ALGEBRAIC LEMMAS

Given a primitive N-th root of unity with N odd, let $\mathcal{W}_q$ be the algebra defined by generators $U^{\pm 1}$, $V^{\pm 1}$, $W^{\pm 1}$ and by the relations $VU = q^4UV$, $WV = q^4VW$, $UW = q^4WU$. It is isomorphic to the Chekhov-Fock algebra $\mathcal{T}^q_\lambda$ for any ideal triangulation λ of the 1-puncture torus. The central element is $H = q^2UVW$. This will not cause any trouble because we are going to deal with automorphisms of $\mathcal{W}$ instead of isomorphisms $\Phi^q_{\lambda\lambda'}$ between different algebras.

Lemma 6 (See for instance [1, Sect. 4])**.** *Every finite-dimensional irreducible representation of $\mathcal{W}_q$ is conjugate to $\chi_{u,v,h} : \mathcal{W}_q \to \mathrm{End}(\mathbb{C}^N)$, defined by*

$$\chi_{u,v,h}(U) = u \begin{pmatrix} 1 & 0 & 0 & \cdots & 0 \\ 0 & q^4 & 0 & \cdots & 0 \\ 0 & 0 & q^8 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & q^{4(N-1)} \end{pmatrix}, \quad \chi_{u,v,h}(V) = v \begin{pmatrix} 0 & 1 & 0 & \cdots & 0 \\ 0 & 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & 1 \\ 1 & 0 & 0 & \cdots & 0 \end{pmatrix},$$

and

$$\chi_{u,v,h}(W) = \frac{q^{-2}h}{uv} \begin{pmatrix} 0 & 0 & 0 & \cdots & q^{-4(N-1)} \\ 1 & 0 & 0 & \cdots & 0 \\ 0 & q^{-4} & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \cdots & q^{-4(N-2)} & 0 \end{pmatrix}.$$

Note that in this case $\chi_{u,v,h}(H) = h \,\mathrm{id}_{\mathbb{C}^N}$. □

Let $\widehat{\mathcal{W}}_q$ be the fraction division algebra of $\mathcal{W}_q$. Consider the automorphism $\mathcal{R} : \widehat{\mathcal{W}}_q \to \widehat{\mathcal{W}}_q$ defined by the property that

$$\begin{aligned} \mathcal{R}(U) &= \big(1+qU^{-1}\big)^{-1}\big(1+q^3U^{-1}\big)^{-1}W \\ \mathcal{R}(V) &= \big(1+qU\big)\big(1+q^3U\big)V \\ \mathcal{R}(W) &= U^{-1}. \end{aligned} \tag{4.1}$$

One easily see that $\mathcal{R}$ does define an algebra isomorphism, namely that $\mathcal{R}(U)$, $\mathcal{R}(V)$ and $\mathcal{R}(W)$ satisfy the same relations as U, V and W (compare Proposition 5 in [5]), and that $\mathcal{R}$ is invertible.

Lemma 7. *If* $\chi_{u,v,h}$ *is a standard representation as in Lemma 6, then*

$$\chi_{u,v,h} \circ \mathcal{R}(X) = C_R \cdot \chi_{u',v',h'}(X) \cdot C_R^{-1}, \qquad \forall\ X \in \mathcal{W}_q,$$

where

$$(u')^N = \frac{h^N}{u^N v^N (1+u^{-N})^2}, \qquad (v')^N = (1+u^N)^2\, v^N, \qquad h' = h$$

and where the matrix $C_R(u,v,u',v',h) \in \mathrm{GL}_N(\mathbb{C})$ *is defined by its entries*

$$(C_R)_{ij} = q^{2(j-i)^2}\ \Big(\frac{u'v'}{uh}\Big)^{j-i}\ \Big(\frac{v'}{v}\Big)^i \prod_{\alpha=1}^{i} \frac{1}{\big(1+q^{4\alpha-3}u\big)\big(1+q^{4\alpha-1}u\big)}.$$

Proof. Lemma 5 shows that $\chi_{u,v,h} \circ \mathcal{R}$ is isomorphic to a standard representation $\chi_{u',v',h'}$ with $(u')^N = \frac{h^N}{u^N v^N (1+u^{-N})^2}$, $(v')^N = (1+u^N)^2\, v^N$ and $h' = h$. To compute the conjugation matrix C_R, the last two equalities of (4.1) provide

$$\begin{aligned} (C_R)_{ij} &= \frac{v'}{v}\, \frac{1}{(1+q^{4i-3}u)(1+q^{4i-1}u)}\, (C_R)_{i-1\ j-1} \\ (C_R)_{ij} &= q^{-4i}\ q^{4(j-1)}\ \frac{u'v'}{uq^{-2}h}\ (C_R)_{i\ j-1} \end{aligned}$$

and these inductive relations immediately give the entries of C_R. □

Similarly, let the isomorphism $\mathcal{L} : \widehat{\mathcal{W}}_q \to \widehat{\mathcal{W}}_q$ be defined by

$$\begin{aligned} \mathcal{L}(U) &= \big(1+qV^{-1}\big)^{-1}\big(1+q^3V^{-1}\big)^{-1}U \\ \mathcal{L}(V) &= \big(1+qV\big)\big(1+q^3V\big)W \\ \mathcal{L}(W) &= V^{-1}, \end{aligned} \tag{4.2}$$

Lemma 8. *If* $\chi_{u,v,h}$ *is a standard representation as in Lemma 6, then*

$$\chi_{u,v,h} \circ \mathcal{L}(X) = C_L \cdot \chi_{u'',v'',h''}(X) \cdot C_L^{-1}, \qquad \forall\ X \in \mathcal{W}_q,$$

where

$$(u'')^N = \frac{u^N}{(1+v^{-N})^2}, \qquad (v'')^N = \frac{h^N(1+v^N)^2}{u^N v^N}, \qquad h'' = h$$

where $C_L = G\widetilde{C}_L$ *is the product of matrices* G, $\widetilde{C}_L(u,v,u'',v'',h) \in \mathrm{GL}_N(\mathbb{C})$ *with entries* $G_{ij} = q^{4ij}$ *and*

$$(\widetilde{C}_L)_{ij} = q^{2(j-i)^2+2i^2}\ \Big(\frac{u''v''}{vh}\Big)^{j-i}\ \Big(\frac{uvv''}{h}\Big)^i \prod_{\alpha=1}^{i} \frac{1}{\big(1+q^{4\alpha-3}v\big)\big(1+q^{4\alpha-1}v\big)}.$$

Proof. The proof is similar to the proof of the previous lemma, except that $\chi_{u,v,h}(V)$ is not diagonal. This inconvenience is bypassed in the following way. Define another irreducible representation $\mu_{u,v,h}$ by

$$\mu_{u,v,h}(U) = u \begin{pmatrix} 0 & 0 & 0 & \cdots & 1 \\ 1 & 0 & 0 & \cdots & 0 \\ 0 & 1 & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \cdots & 1 & 0 \end{pmatrix}, \quad \mu_{u,v,h}(V) = v \begin{pmatrix} 1 & 0 & 0 & \cdots & 0 \\ 0 & q^4 & 0 & \cdots & 0 \\ 0 & 0 & q^8 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & q^{4(N-1)} \end{pmatrix},$$

$$\mu_{u,v,h}(W) = \frac{q^{-2}h}{uv} \begin{pmatrix} 0 & 1 & 0 & \cdots & 0 \\ 0 & 0 & q^{-4} & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & q^{-4(N-2)} \\ q^{-4(N-1)} & 0 & 0 & \cdots & 0 \end{pmatrix}.$$

One easily computes that $G \cdot \mu_{u,v,h}(X) \cdot G^{-1} = \chi_{u,v,h}(X)$, with $G_{ij} = q^{4ij}$. Let $\widetilde{C}_L$ be the matrix such that $\mu_{u,v,h} \circ \mathfrak{L}(X) = \widetilde{C}_L \cdot \chi_{u'',v'',h}(X) \cdot \widetilde{C}_L^{-1}$. Obviously $C_L = G\widetilde{C}_L$. As in the previous lemma, we obtain the inductive equations

$$(\widetilde{C}_L)_{ij} = q^{4(i-1)} \left(\frac{uvv''}{q^{-2}h}\right) \frac{1}{\left(1+q^{4i-3}v\right)\left(1+q^{4i-1}v\right)} (\widetilde{C}_L)_{i-1\ j-1}$$

$$(\widetilde{C}_L)_{ij} = q^{-4i}\, q^{4(j-1)}\, \frac{u''v''}{vq^{-2}h}\, (\widetilde{C}_L)_{i\ j-1}$$

which provide the entries of $\widetilde{C}_L$. □

Remark 9. Note that, although the quantities u, v, h, u', v', involved in the definition of the matrix C_R are related by equations involving only their N–th powers, the matrix itself depends on the choice of these N–roots. The same holds for C_L. In the next section, the N–powers will be determined by geometric information but the actual computation will depend on actual choices of N–roots for this geometric data.

5. The invariant in the case of the 1–puncture torus

We now compute the invariant C_φ of Theorem 3 in the case of the 1-puncture torus T. The computation is here greatly simplified by the fact that the mapping class group of T is isomorphic to $\mathrm{SL}_2(\mathbb{Z})$. In addition, recall that a mapping class is pseudo-Anosov exactly when the corresponding element of $\mathrm{SL}_2(\mathbb{Z})$ has a trace of absolute value greater than 2.

Proposition 10. *Let A be an element of $\mathrm{SL}_2(\mathbb{Z})$ with $|\mathrm{Tr}(A)| > 2$. Then the conjugacy class of A in $\mathrm{SL}_2(\mathbb{Z})$ contains an element of the form*

$$\begin{pmatrix} 1 & a_1 \\ 0 & 1 \end{pmatrix}\begin{pmatrix} 1 & 0 \\ b_1 & 1 \end{pmatrix}\begin{pmatrix} 1 & a_2 \\ 0 & 1 \end{pmatrix}\begin{pmatrix} 1 & 0 \\ b_2 & 1 \end{pmatrix} \cdots \begin{pmatrix} 1 & a_n \\ 0 & 1 \end{pmatrix}\begin{pmatrix} 1 & 0 \\ b_n & 1 \end{pmatrix}$$

where $n > 0$ and the a_i, b_i are positive integers. Moreover, the right hand side is unique up to cyclic permutation of the factors $\begin{pmatrix} 1 & a_i \\ 0 & 1 \end{pmatrix}$ *and* $\begin{pmatrix} 1 & 0 \\ b_i & 1 \end{pmatrix}$. □

In other words, after conjugation, every mapping class of T can be represented by a matrix $A = A_1A_2\cdots A_n$ with $A_i = R$ or L , where $L = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ and $R = \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}$.

The identification between $H_1(T) \cong \mathbb{Z}^2$ assigns a slope in $\mathbb{Q} \cup \{\infty\}$ to each edge of an ideal triangulation λ of T. Consider the ideal triangulation $\lambda_{(0)}$ whose edges have respective slopes 0, ∞ and 1. Then define ideal triangulations $\lambda_{(0)}$, $\lambda_{(1)}$, …, $\lambda_{(n)} = \varphi(\lambda_{(0)})$ by the property that $\lambda_{(i)} = A_1A_2\dots A_i(\lambda_{(0)})$.

It is immediate that the edges $\lambda_{(i)1}$, $\lambda_{(i)2}$ and $\lambda_{(i)3}$ of $\lambda_{(i)}$ all have non-negative slope. Choose the indexing of these edges so that $\lambda_{(i)1}$ has the lowest slope and $\lambda_{(i)2}$ has the highest slope of the three. Note that with this convention the edges $\lambda_{(i)1}$, $\lambda_{(i)2}$ and $\lambda_{(i)3}$ occur counterclockwise in this order around the two triangles of $\lambda_{(i)}$.

The key observation is the following:

- if $A_i = R$, then $\lambda_{(i)}$ is obtained from $\lambda_{(i-1)}$ by a diagonal exchange along the edge $\lambda_{(i-1)1}$, followed by a reindexing exchanging $\lambda_{(i)1}$ and $\lambda_{(i)3}$;
- if $A_i = L$, then $\lambda_{(i)}$ is obtained from $\lambda_{(i-1)}$ by a diagonal exchange along the edge $\lambda_{(i-1)2}$, followed by a reindexing exchanging $\lambda_{(i)2}$ and $\lambda_{(i)3}$.

This property is illustrated in Figure 3 for the case where $i = 1$, and the other cases follow from this one by observing that

$$\lambda_{(i)} = (A_1A_2\dots A_{i-1})A_i(A_1A_2\dots A_{i-1})^{-1}(\lambda_{(i-1)}).$$

Incidentally, this accounts for the somewhat unnatural order of the A_k in the definition of $\lambda_{(i)}$.



Figure 3.

For the complete hyperbolic metric on the mapping torus $M_\varphi = S \times [0,1]/\sim$ with its complete hyperbolic metric, each ideal triangulation $\lambda_{(i)}$ determines a unique pleated surface $f_{\lambda_{(i)}} : T \to M_\varphi$ with pleating locus $\lambda_{(i)}$. The geometry of this pleated surface associates complex weights $x_{(i)1}$, $x_{(i)2}$, $x_{(i)3} \in \mathbb{C} - \{0\}$ to the edges of $\lambda_{(i)}$, corresponding to the exponential shear-bend coordinates of $f_{\lambda_{(i)}}$ along the components of its pleating locus.

Remark 11. In [3], Guéritaud proves the remarkable fact that the pleated surfaces $f_{\lambda_{(i)}}$ are embeddings. In addition, one passes from$f_{\lambda_{(i-1)}}$ to $f_{\lambda_{(i)}}$ by a diagonal exchange across a positively oriented hyperbolic ideal tetrahedra with dihedral angles strictly between 0 and π.

Theorem 12. *Given any pseudo-Anosov diffeomorphism $\varphi : T \to T$, if φ has the canonical decomposition $A_1 A_2 \cdots A_n$ with $A_i = R$ or L as in Proposition 10, then the conjugacy class of C_φ has a matrix representative*

$$C_\varphi = C_1 C_2 \cdots C_n$$

where

- *if $A_i = R$, then $C_i = C_R(u_{i-1}, v_{i-1}, u_i, v_i, 1)$, defined in Lemma 7, with*

$$(u_i)^N = \frac{1}{u_{i-1}^N v_{i-1}^N (1 + u_{i-1}^{-N})^2}, \qquad (v_i)^N = (1 + u_{i-1}^N)^2\, v_{i-1}^N,$$

- *if $A_i = L$, then $C_i = C_L(u_{i-1}, v_{i-1}, u_i, v_i, 1)$, defined in Lemma 8, with*

$$(u_i)^N = \frac{u_{i-1}^N}{(1 + v_{i-1}^{-N})^2}, \qquad (v_i)^N = \frac{(1 + v_{i-1}^N)^2}{u_{i-1}^N v_{i-1}^N},$$

and, the initial values u_0, v_0, h are chosen so that

$$u_n = u_0, \qquad v_n = v_0, \qquad h = 1.$$

Proof. The action of the word $A_1 A_2 \cdots A_n$ produces a sequence of ideal triangulations

$$\lambda = \lambda_{(0)} \xrightarrow{A_1} \lambda_{(1)} \xrightarrow{A_2} \lambda_{(2)} \longrightarrow \cdots \xrightarrow{A_n} \lambda_{(n)} = \varphi(\lambda),$$

which in turn gives a composition of isomorphisms

$$\widehat{\mathcal{T}}^q_{\varphi(\lambda)} \xrightarrow{\Phi^q_{\lambda_{(n-1)}\lambda_{(n)}}} \widehat{\mathcal{T}}^q_{\lambda_{(n-1)}} \longrightarrow \cdots \xrightarrow{\Phi^q_{\lambda_{(0)}\lambda_{(1)}}} \widehat{\mathcal{T}}^q_{\lambda} .$$

By sending $X_{\lambda_{(i)1}}$, $X_{\lambda_{(i)2}}$, $X_{\lambda_{(i)3}}$ respectively to U, V, W we identify each $\widehat{\mathcal{T}}^q_{\lambda_{(i)}}$ to the algebra $\widehat{\mathcal{W}}_q$ of §4. Then we get an automorphism $\mathcal{A}_1 \mathcal{A}_2 \cdots \mathcal{A}_n : \widehat{\mathcal{W}}_q \to \widehat{\mathcal{W}}_q$ with $\mathcal{A}_i \in \{\mathcal{R}, \mathcal{L}\}$. This automorphism is conjugate to the automorphism $\Phi^{-1} \circ \Phi^q_{\lambda\varphi(\lambda)} : \widehat{\mathcal{T}}^q_{\varphi(\lambda)} \to \widehat{\mathcal{T}}^q_{\varphi(\lambda)}$, ($\Phi : \widehat{\mathcal{T}}^q_{\varphi(\lambda)} \to \widehat{\mathcal{T}}^q_{\lambda}$ is the natural identification). If we choose a standard representation $\chi_{u,v,h}$, then Lemma 7 and Lemma 8 tell us that for any i, $1 \le i \le n$, we have

$$\begin{aligned}&\chi_{u,v,h} \circ \mathcal{A}_1\mathcal{A}_2 \cdots \mathcal{A}_i(X)\\ =&C_1(u_0, v_0, u_1, v_1, h) \cdot \Big(\chi_{u_1,v_1,h} \circ \mathcal{A}_2 \cdots \mathcal{A}_i(X)\Big) \cdot C_1(u_0, v_0, u_1, v_1, h)^{-1}\\ =&\cdots\\ =&(C_1 \cdots C_i)\Big(\chi_{u_k,v_k,h}(X)\Big)(C_1 \cdots C_i)^{-1}\end{aligned}$$

for all $X \in \mathcal{W}_q$. In particular, the hyperbolic metric of the mapping torus M_φ provides a interesting initial standard representation χ_{u_0,v_0,h_0} such that $u_n^N = u_0^N$, $v_n^N = v_0^N$ (We have interpreted u_i^N, v_i^N to be the shear-bend coordinates $x_{(i)1}$, $x_{(i)2}$). We choose the standard representations from their conjugacy classes so that $u_n = u_0$, $v_n = v_0$, and we choose $h = 1$ because it is natural to work on the "cusped quantum Teichmüller space" (see [1, 5]) in which we should put the "cusp condition" $H^2 = 1$, and a geometric consideration fixes $H = 1$. The above equation applied to $i = n$ gives

$$\begin{aligned}&\chi_{u_0,v_0,h_0} \circ \mathcal{A}_1\mathcal{A}_2 \cdots \mathcal{A}_n(X)\\ =&(C_1 \cdots C_n)\Big(\chi_{u_0,v_0,h_0}(X)\Big)(C_1 \cdots C_n)^{-1}\end{aligned}$$

Comparing with equation 2.1, the product of matrices $C_1 \cdots C_n$ turns out to be a representative of the invariant C_φ.

□

6. The invariant in the case of the 4-puncture sphere

We identify the 4-puncture sphere to be the quotient $S \cong (\mathbb{R}^2 - \mathbb{Z}^2)/G$ where G is the group generated by four elements $(x, y) \mapsto (-x, y)$, $(x, y) \mapsto (x+2, y)$, $(x, y) \mapsto (x, -y)$ and $(x, y) \mapsto (x, y+2)$. The mapping class group is the semi-directed product of $\mathrm{PSL}_2(\mathbb{Z})$ on $\mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}/2\mathbb{Z}$. The image of a diffeomorphism φ can be represented by a matrix $A_\varphi \in \mathrm{SL}_2(\mathbb{Z})$.



Figure 4. An ideal triangulation λ of S.

We will restrict attention to ideal triangulation of S which are isomorphic to the 2-skeleton of an ideal tetrahedron, as in the figure above. This is equivalent to the property that each puncture is adjacent to exactly 3 edges.

If $\varphi : S \to S$ is associated to the matrix $A_\varphi \in \mathrm{S}L_2(\mathbb{Z})$, then φ is pseudo-Anosov exactly when $|\mathrm{Tr}\,\varphi| > 2$. Similar to the 1-puncture torus, such a matrix has an LR decomposition. The difference now is that, to stay within tetrahedral ideal triangulations, the L and R are now associated to the product of two diagonal exchanges along opposite edges. More precisely, with the indexing convention of the figure above, L corresponds to $\Delta_2\Delta_4$ and R corresponds to $\Delta_1\Delta_3$. By [5], the Chekhov-Fock algebra $\mathcal{T}_\lambda^q$ (λ as in the figure) is generated by X_1, X_2 and the central elements

$$P_1 = qX_1X_2X_5, \qquad P_2 = q^{-1}X_2X_3X_6, \qquad P_3 = qX_3X_4X_5, \qquad P_4 = qX_1X_4X_6,$$
$$H = q^2X_1X_2X_3X_4X_5X_6.$$

There exist automorphisms $\mathcal{R}$ and $\mathcal{L}$ of the Chekhov-Fock algebra $\mathcal{T}_\lambda^q$ which fix the central elements and map X_1, X_2 as follows,

$$\begin{aligned}\mathcal{R}(X_1) &= \big(1 + qX_1^{-1}\big)^{-1}\big(1 + qX_3^{-1}\big)^{-1}X_6,\\ \mathcal{R}(X_2) &= (1 + qX_1)(1 + qX_3)X_2,\end{aligned}$$

and

$$\begin{aligned}\mathcal{L}(X_1) &= \big(1+qX_2^{-1}\big)^{-1}\big(1+qX_4^{-1}\big)^{-1}X_1,\\ \mathcal{L}(X_2) &= (1+qX_2)(1+qX_4)X_5.\end{aligned}$$

If we use the standard representation

$$\chi_{u,v,h,p}(X_1) = u\begin{pmatrix} 1 & 0 & 0 & \cdots & 0\\ 0 & q^2 & 0 & \cdots & 0\\ 0 & 0 & q^4 & \cdots & 0\\ \vdots & \vdots & \vdots & \ddots & \vdots\\ 0 & 0 & 0 & \cdots & q^{2(N-1)}\end{pmatrix}, \quad \chi_{u,v,h,p}(X_2) = v\begin{pmatrix} 0 & 1 & 0 & \cdots & 0\\ 0 & 0 & 1 & \cdots & 0\\ \vdots & \vdots & \vdots & \ddots & \vdots\\ 0 & 0 & 0 & \cdots & 1\\ 1 & 0 & 0 & \cdots & 0\end{pmatrix},$$

and

$$\chi_{u,v,h,p}(H) = h\ \mathrm{id}_{\mathbb{C}^N}, \quad \chi_{u,v,h,p}(P_j) = p_j\ \mathrm{id}_{\mathbb{C}^N}, \quad j = 1,2,3,4,$$

then we have similar algebraic Lemmas for the 4-puncture sphere, with the new matrix functions,

$$(C_R^*)_{ij} = q^{(i-j)(i-j-2)}\left(\frac{v'}{v}\right)^j (uvp_3u')^{j-i}\prod_{\alpha=1}^{i-j}\frac{1}{(1+q^{1-2\alpha}u^{-1})\left(1+\frac{q^{-1-2\alpha}h}{up_2p_3}\right)} \quad\times$$

$$\times \quad \prod_{\beta=i-j+1}^{i}\frac{1}{(1+q^{2\beta-1}u)\left(1+\frac{q^{2\beta+1}up_2p_3}{h}\right)}$$

$$G_{ij}^* = \ q^{2ij}$$

$$(\widetilde{C}_L^*)_{ij} = q^{j^2}\left(\frac{u'}{u}\right)^{j-i}\left(\frac{uvv'}{p_1}\right)^j \qquad \prod_{\alpha=1}^{i-j}\frac{1}{(1+q^{1-2\alpha}v^{-1})\left(1+\frac{q^{1-2\alpha}h}{vp_3p_4}\right)} \quad\times$$

$$\times \quad \prod_{\beta=i-j+1}^{i}\frac{1}{(1+q^{2\beta-1}v)\left(1+\frac{q^{2\beta-1}vp_3p_4}{h}\right)}$$

The same arguments as in §5 then give:

Theorem 13. *Given any pseudo-Anosov diffeomorphism $\varphi : S \to S$, if φ has the canonical decomposition $A_1A_2\cdots A_n$ with $A_i = R$ or L as in Proposition 10, then the conjugacy class of C_φ has a matrix representative*

$$C_\varphi^* = C_1^*C_2^*\cdots C_n^*$$

where

- *if $A_i = R$, then $C_i^* = C_R^*(u_{i-1}, v_{i-1}, u_i, v_i)$, defined above, with*

$$(u_i)^N = -\frac{1}{u_{i-1}^N v_{i-1}^N (1+u_{i-1}^{-N})^2}, \qquad (v_i)^N = (1+u_{i-1}^N)^2\, v_{i-1}^N,$$

- *if $A_i = L$, then $C_i^* = C_L^*(u_{i-1}, v_{i-1}, u_i, v_i, 1) = G^*\cdot\widetilde{C}_L^*(u_{i-1}, v_{i-1}, u_i, v_i, 1)$, defined above, with*

$$(u_i)^N = \frac{u_{i-1}^N}{(1+v_{i-1}^{-N})^2}, \qquad (v_i)^N = -\frac{(1+v_{i-1}^N)^2}{u_{i-1}^N v_{i-1}^N},$$

and, the initial values u_0, v_0, h, p are chosen so that

$$u_n = u_0, \qquad v_n = v_0, \qquad h = 1, \qquad p_j = 1,\ \forall\ j.$$

□

REFERENCES

[1] Francis Bonahon, Xiaobo Liu, *Representations of the quantum Teichmüller space and invariants of surface diffeomorphisms*, Geom. Topol. 11 (2007), 889–937.

[2] Vladimir V. Fock, Leonid O. Chekhov, *Quantum Teichmüller spaces* (Russian) Teoret. Mat. Fiz. 120 (1999), 511–528; translation in Theoret. and Math. Phys. 120 (1999), 1245–1259.

[3] François Guéritaud, *On canonical triangulations of once-punctured torus bundles and two-bridge link complements*, with an appendix by David Futer, Geom. Topol. 10 (2006), 1239–1284 (electronic).

[4] Rinat Kashaev, *Quantization of Teichmüller spaces and the quantum dilogarithm*, Lett. Math. Phys. 43 (1998), 105–115.

[5] Xiaobo Liu, *The quantum Teichmüller space as a noncommutative algebraic object*, `arXiv:math.GT/0408361`.

DEPARTMENT OF MATHEMATICS, COLUMBIA UNIVERSITY, NEW YORK, NY 10027, U.S.A.
E-mail address: `xiaoboli@math.columbia.edu`